

Solitary Solutions fo Nonlinear Differential-difference Equations via Adomain Decomposition Method

M A Abdou *

Theoretical Research Group,Physics Department, Faculty of Science,Mansoura University,35516 Mansoura,Egypt

(Received 22 June 2010 , accepted 16 March 2011)

Abstract: Here,Adomian decomposition (ADM) method has been used for finding approximate and numerical solutions of nonlinear differential difference equations arising in mathematical physics.Two models of special interst in physics,namely, the Hybrid nonlinear differential difference equation and Relativistic Toda coupled nonlinear differential-difference equation are chosen to illustrate the validity and the great potential of the proposed method.Comparisons are made between the results of the proposed method and exact solutions. It is worthwhile to mention that the Adomian decomposition method is also easy to be applied to other nonlinear differential difference equation arising in physics.

Keywords: Hybrid nonlinear difference equation, Relativistic Toda coupled nonlinear difference equation,Adomian Decomposition method

1 Introduction

The nonlinear differential-difference equations (DDEs) have been the focus of many nonlinear studies.The DDEs play an important role in modelling complicated physical phenomena such as particle vibrations in lattices,currents ow in electrical networks,and pulses in biological chains.The solutions of these DDEs can provide numerical simulations of nonlinear partial di erential equations, queuing problems, and discretizations in solid state and quantum physics.Since the works of Fermi et al. in the 1950s [1],there were quite a number of research works developed during the last decades on DDEs.For instances,Levi and his coworkers analyzed the condition for existence of higher symmetries for a class of DDEs [2, 3],Yamilov and his co-workers [4, 5] ade outstanding contribution to the classification of DDEs,integrability tests and connections between integrable PDEs and DDEs [6 – 23]

The Adomian Decomposition (ADM) Method [24 – 30] has been used to solve effectively,easily,and accurately a large class of linear and nonlinear equations,solutions partial, deterministic or stochastic differential equations with approximates which converge rapidly.

The paper has been organized as follows.**In Section 2**,we extended the ADMto solving nonlinear differential-difference equations with initial condition,an brief outline of the methodology is presented.**In Section 3**,two models arising in physics are chosen to illustrate the validity of the Adomian decomposition method in solving NDDE(s).Finally, discussions and conclusions are presented in **Section 4**.

2 Methods and its Applications

For a given the nonlinear differential-difference equation as

$$L(u_n) + R(u_n, u_{n+1}, u_{n-1} + \dots) + N(u_n, u_{n+1}, u_{n-1}, \dots) = g, \quad (1)$$

with prescribed conditions,where u_n is the unknown function, L is the highest order derivative which is assumed to be n easily invertible, R is a linear differential operator of less order than L , $N(u_n, u_{n+1}, u_{n-1}, \dots)$ represents the nonlinear

* E-mail address: m.abdou.eg@yahoo.com

term and g_n is the source term. Assuming the inverse operator L^{-1} exists and it can be taken as the definite integral with respect to from t_0 to t , i.e.

$$L^{-1} = \int_{t_0}^t (\cdot) dt \quad (2)$$

Applying L^{-1} to both sides of (1) with initial conditions, we have

$$u_n = f(x) - L^{-1}[R(u_n, u_{n+1}, u_{n-1} + \dots) + N(u_n, u_{n+1}, u_{n-1}, \dots)], \quad (3)$$

where the function $f(x)$ represents the term arising from integrating the source term g and from using the given initial or boundary conditions, all are assumed to be prescribed.

The Adomian decomposition, assumes a series that the unknown function $u_n(t)$ can be expressed by an infinite series as

$$u_n(t) = \sum_{m=0}^{\infty} u_{m,n}(t) \quad (4)$$

The nonlinear operator $N(u_n, u_{n+1}, u_{n-1}, \dots)$ can be decomposed by an infinite series of polynomials given by

$$N(u_n, u_{n+1}, u_{n-1}, \dots) = \sum_{m=0}^{\infty} A_{m,n}, \quad (5)$$

where $A_{m,n}$ are the Adomian's polynomials. To determine the Adomian polynomials, we introduce a parameter λ and (5) becomes

$$N\left(\sum_{m=0}^{\infty} u_{m,n}(t)\lambda^m, \sum_{m=0}^{\infty} u_{m,n+1}(t)\lambda^m, \sum_{m=0}^{\infty} u_{m,n-1}(t)\lambda^m, \dots\right) = \sum_{m=0}^{\infty} A_{m,n}\lambda^m, \quad (6)$$

$$A_{m,n} = \frac{1}{m!} \left[\frac{d^m}{d\lambda^m} N\left(\sum_{m=0}^{\infty} u_{m,n}(t)\lambda^m, \sum_{m=0}^{\infty} u_{m,n+1}(t)\lambda^m, \sum_{m=0}^{\infty} u_{m,n-1}(t)\lambda^m, \dots\right) \right]_{\lambda=0}, \quad (7)$$

This formula is easy to compute by using Maple software or by setting a computer code to get as many polynomials as we need in the calculation of the numerical as well as explicit solutions. Knowing the zero component $u_n(0)$, the remaining components where $n > 1$ can be determined by using recurrence relation

$$u_0(n) = f(x), \quad (8)$$

$$u_m(n+1, t) = -L^{-1}[R(u_{m,n}, u_{m,n+1}, u_{m,n-1} + \dots) + A_{m,n}], \quad m > 0 \quad (9)$$

For the numerical computation, we consider the expression as follows

$$\phi_{n,k} = \sum_{m=0}^k u_{m,n}, \quad (10)$$

denotes the n -term approximation to $u_n(t)$.

3 Applications

To illustrate the effectiveness and the advantages of the proposed method. Here, we consider two models of nonlinear differential-difference equations of special interest physically, namely, the Hybrid nonlinear differential difference equation and Relativistic Toda coupled nonlinear differential-difference equation.

3.1 Example(1).The Hybrid nonlinear differential difference equation

Let us first consider the Hybrid nonlinear differential difference equation [20]

$$\frac{\partial u_n}{\partial t} = (\alpha - u_n^2)(u_{n+1} - u_{n-1}), \tag{11}$$

subject to the initial condition

$$u_n(0) = \sqrt{-\alpha} \tanh(d/2)(1 + \cosh(d)) \operatorname{sech}(dn - 2), \tag{12}$$

where d is constant. The Hybrid nonlinear difference Eq.(11) describe the discretization of the KdV and modified KdV equations. Applying ADM, Eq.(11) can be rewritten in the operator form

$$L[u_n(t)] = [R(u) + N(u)], \tag{13}$$

where $L = \frac{\partial}{\partial t}$ is a linear operator $R(u) = \alpha[u(n + 1) - u(n - 1)]$ is the remainder of the linear operator. The nonlinear term is given by $N(u) = -u^2(n)(u(n + 1) - u(n - 1))$. Operating with L^{-1} on to both sides of Eq.(13) with initial condition (12) gives

$$u_n(t) = u_n(0) + L^{-1}[R(u) + N(u)] \tag{14}$$

The Adomian decomposition method assumes an infinte series solutions of the unknown function $u_n(t)$ in the form

$$u_n(t) = \sum_{m=0}^{\infty} u_m(n, t) \tag{15}$$

It is to be noted that the subscript m does not stand for the m th lattice any more, it means the m th element is the decomposition series. Similarly, the nonlinear operator $N(u)$ ca be decomposed as

$$N(u) = \sum_{m=0}^{\infty} A_m(u_0, u_1, u_2, \dots, u_m), \tag{16}$$

where A_m are the appropriate Adomian polynomials, which is defined as [24, 25]. The first four components od A_m are given by [21].

According to Eqs.(14),(15) and (16), the first few components of the decomposition series $u_m(n, t)$ are readily found to be

$$u_0(n) = \sqrt{-\alpha} \tanh(d/2)(1 + \cosh(d)) \operatorname{sech}(dn - 2),$$

$$u_1(n, t) = L^{-1}[\alpha[u_0(n + 1, t) - u_0(n - 1, t)] + A_0],$$

$$u_2(n, t) = L^{-1}[\alpha[u_1(n + 1, t) - u_1(n - 1, t)] + A_1],$$

$$u_3(n, t) = L^{-1}[\alpha[u_2(n + 1, t) - u_2(n - 1, t)] + A_2],$$

$$u_{m+1}(n, t) = L^{-1}[\alpha[u_m(n + 1, t) - u_m(n - 1, t)] + A_m], \tag{17}$$

and so on. The rest of the components of $u_m(n, t)$ can be directly evaluated via (17) in a similar way. It is worth noting that the accuracy of the Adomian approach is significantly enhanced by calculation as many more terms as we like. This may be made but only at the expense of a considerable increase in the complexity of analysis. The Explicit forms for u_1, u_2, u_3, \dots are obtained and for simplicity should be omitted here.

To verify numerically whether the proposed methodology lead to high accuracy, we evaluate the numerical solutions using the fourth-term approximation and compared it with the exact analytical solution. So, we could get fourth-order approximation $\phi_5(n) = \sum_{m=0}^5 u_m(n, t)$. Figs.[1.(a,b)] shows comparisons between the numerical approximate solutions obtained by ADM with a fixed values of d, t for a different values of n and exact solutions [20], which proofs the two solutions are quite good. We could say that solution to the Hybrid nonlinear differential difference equation obtained by ADM agree well with the exact solutions [20]

$$u_n(t) = \sqrt{-\alpha} \tanh(d/2)(1 + \cosh(d)) \operatorname{sech}[dn - 2\alpha \tanh(d/2)(1 + \cosh d)t - 2], \tag{18}$$

3.2 Example(2).Relativistic Toda coupled nonlinear differential-difference equation

A second instructive model is the Relativistic Toda coupled nonlinear differential difference equation [20]

$$\frac{\partial u_n(t)}{\partial t} = (1 + \alpha u_n(t))(v_n(t) - v_{n-1}(t)), \quad (19)$$

$$\frac{\partial v_n(t)}{\partial t} = v_n(t)(u_{n+1}(t) - u_n(t) + \alpha v_{n+1}(t) - \alpha v_{n-1}(t)), \quad (20)$$

subject to the initial conditions

$$u_n(0) = -1 - c_1 \coth(d_1) + c_1 \tanh[d_1 n], \quad (21)$$

$$v_n(0) = c_1 \coth(d_1) - c_1 \tanh[d_1 n], \quad (22)$$

where c_1 and d_1 are constants. The Toda lattice difference Eqs.(19) and (20) describe vibrations in mass spring lattices with an exponential interaction force. Applying ADM, Eqs.(19) and (20) can be written in an operator form

$$L[u_n(t)] = [v(n) - v(n-1)] + \alpha M[u(n), v(n), v(n-1)], \quad (23)$$

$$L[v_n(t)] = N[u(n), u(n+1), v(n), v(n+1), v(n-1)], \quad (24)$$

Operating with L^{-1} on to both sides of Eqs.(23) and (24) with initial conditions (21) and (22) gives

$$u_n(t) = u_n(0) + L^{-1}[v(n) - v(n+1) + \alpha M[(u(n), v(n), v(n-1))], \quad (25)$$

$$v_n(t) = v_n(0) + L^{-1}[N(u(n), u(n+1), v(n), u(n+1))] \quad (26)$$

The nonlinear operators $M[u(n), v(n), v(n-1)] = u(n)(v(n) - v(n-1))$ and $N[(u(n), u(n+1), v(n), u(n+1))] = v(n)[u(n+1) - u(n) + \alpha v(n+1) - \alpha v(n-1)]$ are given by the infinite series of Adomian polynomial

$$M(u_n, v_n, v_{n-1}) = \sum_{m=0}^{\infty} A_m, \quad (27)$$

$$N(u_n, u_{n+1}, v_n, v_{n+1}, v_{n-1}) = \sum_{m=0}^{\infty} B_m, \quad (28)$$

where A_m and B_m are the approximate Adomian polynomials which are generated according to algorithms determined

The Adomian decomposition method assumes an infinite series solutions of the unknown functions $u_n(t)$ and $v_n(t)$ in the form

$$u_n(t) = \sum_{m=0}^{\infty} u_m(n, t), \quad (29)$$

$$v_n(t) = \sum_{m=0}^{\infty} v_m(n, t) \quad (30)$$

Inserting Eqs.(29) and (30) into Eqs.(25) and (26) gives

$$u_0(n) = -1 - c_1 \coth(d_1) + c_1 \tanh[d_1 n], \quad (31)$$

$$v_0(n) = c_1 \coth(d_1) - c_1 \tanh[d_1 n],$$

$$u_{m+1}(n, t) = L^{-1}[v_m(n, t) - v_m(n+1, t) + A_m], m > 0$$

$$v_{m+1}(n, t) = L^{-1}[B_m], m > 0 \quad (32)$$

With the aid of the zeroth components of $u_{n,0}$ and $v_{n,0}$, by all terms arise from the initial conditions (21) and (22), and as a result, the remaining components $u_m(n, t)$ and $v_m(n, t)$, $m > 0$ can be determined. The Explicit forms for u_1, u_2, u_3 and v_1, v_2 and v_3 are obtained and for simplicity should be omitted here. To verify numerically whether the proposed methodology lead to high accuracy, we evaluate the numerical solutions using the fourth-term approximation and compared it with the exact analytical solution. So, we could get fourth-order approximation $\phi_5(n) = \sum_{m=0}^5 u_m(n, t)$. Figs. [2.(a-d)] shows comparisons between the numerical approximate solutions obtained by ADM and exact solutions [20], which proofs the two solutions are quite good. It is to be noted that solution to Relativistic Toda coupled nonlinear differential-difference equation obtained by ADM agree well with the exact solutions [20]

$$u_n(t) = -1 - c_1 \coth(d_1) + c_1 \tanh[d_1 n + c_1 t], \quad (33)$$

$$v_n(t) = c_1 \coth(d_1) - c_1 \tanh[d_1 n + c_1 t], \quad (34)$$

where c_1 and d_1 are constants to be determined later.

4 Summary and Discussion

In summary, we successfully apply the Adomian Decomposition method to solve nonlinear NDDEs. As an illustrative example, we employ DDEs-ADM to the Hybrid nonlinear differential difference equation and Relativistic Toda coupled nonlinear differential-difference equation. The comparison are made between the solutions obtained by Adomian Decomposition Method with exact solution. The comparison shows the validity of Adomian Decomposition Method applied to nonlinear differential-difference equation. Since the numerous DDEs in many fields of scientific and the hardness of solving them, this DDEs-ADM will become a much more interesting method to solving nonlinear DDEs in science and engineering.

The results reported here provide further evidence of the usefulness of Adomain decomposition (AMD) method. The ADM was clearly very efficient and powerful technique in finding the solutions of the nonlinear differential difference equations. It is clear that this method avoids linearization and biologically unrealistic assumptions, and provides an efficient numerical solution.

A clear conclusion can be drawn from the numerical results that the ADM algorithm provides highly accurate numerical solutions without spatial discretization for nonlinear differential equations. It is also worth noting that the advantage of the decomposition methodology displays a fast convergence of the solutions.

Finally, we point out that the ADM confirm the correctness of those obtained by other methods. The method is straightforward and concise, and it can also be applied to other nonlinear differential difference equations in mathematical physics. This is our task in future work.


Acknowledgments

The author would like to express sincerely thanks to the referees for useful comments and discussions.


References

- [1] Fermi E, Pasta J, Ulam S. Collected papers of Enrico Fermi. Chicago, IL: Chicago Press; 1965. p. 978.
- [2] Levi D, Ragnisco O. Extension of the spectral transform method for solving nonlinear differential-difference equations. *Lett Nuovo Cimento*, 22(1978):6916.
- [3] Levi D, Yamilov RI. Conditions for the existence of higher symmetries of evolutionary equations on the lattice. *J Math Phys*. 38(1997):664-874.
- [4] Yamilov RI. Construction scheme for discrete Miura transformations. *J Phys A: Gen*. 1994;27:683-951.
- [5] Adler VE, Svinolupov SI, Yamilov RL. Multicomponent Volterra and Toda type integrable equations. *Phys Lett A*, 254(1999):24-36.
- [6] Cherdantsev IYu, Yamilov RI. Master symmetries for differential difference equations of the Volterra type. *Physica D*, 87(1995):14-04.


- [7] Shabat AB, Yamilov RI. Symmetries of nonlinear chains. *Leningrad Math J*, 2(1991):377-400.
- [8] Zhang DJ. Singular solutions in Casoratian form for two differential difference equations. *Chaos, Solitons & Fractals*, 23(2005):133-350.
- [9] Narita K. Soliton solution for a highly nonlinear difference differential equation. *Chaos, Solitons & Fractals*, 3(1993):27983.
- [10] Wang Z, Zhang HQ. New exact solutions to some difference differential equations. *Chin Phys.* 15(2006):22105.
- [11] Wang Z, Zhang HQ. A symbolic computational method for constructing exact solutions to difference differential equations. *Appl.Math.Comput.*,178(2006):431.
- [12] Suris YuB. New integrable systems related to the relativistic Toda lattice. *J Phys A: Math Gen.* 30(1997):1745.
- [13] Suris Yu B. Integrable discretizations for lattice systems: local equations and their Hamiltonian properties. *Rev Math Phys*, 11(1999):727-822.
- [14] Suris YuB. The problem of integrable discretization: Hamiltonian approach. *Progress in mathematics.* Vol.219. Basel: Birkha Verlag; 2003.
- [15] L.Zou,Z.Zong,C.H.Dong, Generalized homotopy analysis method to solve Lotka Voltera equation. *Comut.and Math.with Appl.*56(2008):2289-2293.
- [16] Ahmet, Applying Hes Variational Iteration Method for Solving Differential-Difference Equation, Mathematical Problems in Engineering, 2008, ID 869614, Pag.7.
- [17] Z.Wang,L.Zou, and H.Zhang. Applying homotopy analysis method for solving differential difference equation. *Physics Letters A*, 369(2007):77-84.
- [18] C.Dai and J.Zhang, Jacobian elliptic function method for nonlinear differential-difference equations. *Chaos, Solitons & Fractals*, 27(2006):1042-1047.
- [19] Z Wang, Discrete tanh method for nonlinear difference-differential equations. *Computer Physics Communications*, 10.1016/j.cpc.2009.01.010 in press
- [20] D.Baldwin, U.Goktas and W.Hereman. *Comp.Phys. Commun.*, 162(2004):203-217.
- [21] Lei Wu, Li-dan Xieb, Jie-fang Zhang. Adomian decomposition method for nonlinear differential-difference equations. *Comm.in Non.Sci.and Numer.Simulation*, 14(2009) :12-18.
- [22] Zhen Wang and Hongqing Zhanga, Construct solitary solutions of discrete hybrid equation by Adomian Decomposition. *Method, Chaos, Solitons & Fractals*, 40(2009):676.
- [23] Li Zou, Zhen Wang, Zhi Zong, Generalized Differential transform method to differential difference equation. *Phys.Lett.A*, 373(2009):4142-4151.
- [24] Adomian G. The decomposition method. Boston: Kluwer Acad. Publ.; 1994.
- [25] M A Abdou. Adomian decomposition method for solving the telegraph equation in charged particle transport, *J. Quant. Spectrosc. Radiat. Transfer*, 95 (2005):407-414.
- [26] M A Abdou and A Elhanbaly. Decomposition method for solving a system of coupled fractional-time nonlinear equations. *Physica Scripta*, 73(2006):338-348.
- [27] S A El-Wakil and M A Abdou. New applications of Adomian decomposition method. *Chaos Solitons & Fractals*, 33 (2007):513-522.
- [28] S A El-Wakil, M A Abdou and A Elhanbaly. Adomian decomposition method for solving diffusion-convection reaction equation. *Applied Mathematics and Computation*, 177(2006):729-736.
- [29] A Elhanbaly and M A Abdou, New application of Adomian decomposition method on Fokker-Planck equation. *Applied Mathematics and Computation*, 182(2006):301-312.
- [30] S A El-Wakil, A Elhanbaly and M A Abdou. Adomian decomposition method for solving fractional nonlinear differential equation. *Applied Mathematics and Computation*, 182(2006):313-324.




Fig[1.a].Approximate solution for 5-order approximation of $u_n(t)$ obatned by Adomian decomposition method with a fixed values of $d=0.1, t=15$ values of n of $d=0.1, t=15$ and $\alpha=-0.1$




Fig[1.b].Exact solution of $u_n(t)$ Eq.(18) with a fixed values for a different values of n and $\alpha=-0.1$ for a different




Fig[2.a].Approximate solution for 5-order approximation of a fixed $u_n(t)$ obatned by Adomian decomposition method with values of $d=0.5, t=.1$ and $c=-0.1$ for a different values of n



Fig[2.b].Exact solution of $u_n(t)$ Eq. (32) with a fixed values of $d=0.5, t=.1$ and $c=-0.1$ for a different values of n



Fig[2.c].Approximate solution for 5-order approximation of $v_n(t)$ obatned by Adomian values of $d=0.5, t=.1$ and $c=-0.1$ for a different values of n decomposition method with a fixed



Fig[2.d].Exact solution of $v_n(t)$ with Eq. (33) a fixed values of $d=0.5, t=.1$ and $c=-0.1$ for a different values of n